

RIKEN-TH-173

New Gauged Linear Sigma Models for 8D HyperKähler Manifolds and Calabi-Yau Crystals

YUTAKA BABA^{*} and TA-SHENG TAI[†]

Theoretical Physics Laboratory, RIKEN, Wako, Saitama 351-0198, JAPAN

Abstract

We propose two kinds of gauged linear sigma models whose moduli spaces are real eight-dimensional hyperKähler and Calabi-Yau manifolds, respectively. Here, hyperKähler manifolds have $sp(2)$ holonomy in general and are dual to Type IIB $(p, q)5$ -brane configurations.

On the other hand, Calabi-Yau fourfolds are toric varieties expressed as quotient spaces. Our model involving fourfolds is different from the usual one which is directly related to a symplectic quotient procedure. Remarkable features in newly-found three-dimensional Chern-Simons-matter theories appear here as well, such as *dynamical* Fayet-Iliopoulos parameters, one *dualized photon* and its residual discrete gauge symmetry.

^{*}e-mail address : ybaba@riken.jp

[†]e-mail address : tasheng@riken.jp

1 Introduction

A gauged linear sigma model (GLSM) in two dimensions [1] is capable of describing a curved geometry (typically a symplectic or Kähler quotient space) through gauge theory language. More precisely, the isometry¹ of the transverse internal space gets partially gauged and chiral multiplets on the worldsheet are coupled to corresponding gauge fields. Then, the curved geometry $\mathbb{C}^n//G$ arises as the supersymmetric vacuum (moduli space) of the 2D gauge theory. Note that $G = \mathbb{C}^{\star r}$ stands for r complexified $U(1)$'s with moment maps

$$\mu : \mathbb{C}^n \rightarrow \mathbb{R}^r, \quad \mu^a = \sum_{i=1}^n Q_i^a |\phi_i|^2, \quad a = 1, \dots, r, \quad Q_i^a : \text{charge assignment.} \quad (1.1)$$

ϕ_i parameterizing $\mathbb{C}^n$ represents the lowest component of an $\mathcal{N}=(2,2)$ chiral superfield Φ_i . Also, μ^a 's are called Fayet-Iliopoulos (FI) parameters which bring in r Kähler classes. (1.1) modulo the following gauge symmetry

$$\phi_i \sim \phi_i e^{i \sum_a Q_i^a \lambda^a}, \quad \lambda^a \in \mathbb{R} \quad (1.2)$$

is exactly the vacuum manifold denoted as $\mathbb{C}^n//G$ or $\mu^{-1}(0)/U(1)^r$.

Similarly, a hyperKähler quotient space is defined by $\mathbb{H}^n///\mathbb{G}$ ($\mathbb{H}^n \cong \mathbb{R}^{4n}$) where $\mathbb{G}$ ($\subset$ triholomorphic isometry) is generated by vector fields K 's with $\mathcal{L}_K g = 0$ (g : metric) and $\mathcal{L}_K I^\alpha = 0$. Three complex structures I^α ($\alpha = 1, 2, 3$) are

$$\omega^\alpha = d\mathcal{P}^\dagger \sigma^\alpha \wedge d\mathcal{P}, \quad \mathcal{P} = (q, \tilde{q}^\dagger)^t, \quad \sigma^\alpha : \text{Pauli matrices} \quad (1.3)$$

for each $\mathbb{H}$ described by $ds^2 = |dq|^2 + |d\tilde{q}|^2$ ($q, \tilde{q} \in \mathbb{C}$) and transform as a triplet under $SU(2)$. Our convention is as follows. A quaternion $\mathcal{Q} = y + \mathbf{y}$ ($\bar{\mathcal{Q}} = y - \mathbf{y}$) with a pure imaginary part $\mathbf{y} = ui + vj + wk$ ($y, u, v, w \in \mathbb{R}$) consists of a pair of complex numbers $(q, \tilde{q})$:

$$\mathcal{Q} = -q + \tilde{q}j, \quad \bar{\mathcal{Q}} = -q^\dagger - j\tilde{q}^\dagger. \quad (1.4)$$

For n quaternions, r moment maps associated with $\mathbb{G}$ under a charge matrix Q_i^a will read

$$\sum_{i=1}^n Q_i^a \mathcal{P}_i^\dagger \boldsymbol{\sigma} \mathcal{P}_i = \boldsymbol{\xi}^a, \quad \mathcal{P}_i = (q_i, \tilde{q}_i^\dagger)^t, \quad a = 1, \dots, r$$

¹For a Kähler quotient space $M//G$, the isometry G is required to be symplectic such that vector fields K 's generating it give $\mathcal{L}_K \omega = 0$ where ω is a symplectic form on the symplectic manifold M .

or

$$\sum_{i=1}^n Q_i^a (|q_i|^2 - |\tilde{q}_i|^2) = \xi^{3a}, \quad \sum_{i=1}^n Q_i^a 2q_i \tilde{q}_i = \xi^{1a} - i\xi^{2a}. \quad (1.5)$$

Here, triplets ξ 's are given level sets. By definition $\mathbb{H}^n // \mathbb{G}$ has real $4(n-r)$ dimensions.

In this note, we propose two kinds of GLSMs which have, respectively, 8D hyperKähler manifolds and Calabi-Yau (CY) 4-folds as their moduli spaces. In the former case, we have followed pioneering works [2, 3, 4]. We extend their $\mathcal{N}=(4,4)$ models to include generic 8D hyperKähler geometries dual to Type IIB $(p,q)5$ -brane configurations [6]. Moreover, taking an infra-red (IR) limit leads to frozen kinetic terms of vector-multiplets. By integrating them out, a nonlinear sigma model (NL σ M) can be realized in the Higgs branch; that is, the quotient space metric is now pulled back onto worldsheet's kinetic term. The latter case is an $\mathcal{N}=(2,2)$ model which provides instead a CY 4-fold at IR. It is not the traditional one [1] which executes a symplectic quotient because it possesses all key features of 3D $\mathcal{N}=2$ Chern-Simons-matter theories on a stack of M2-branes probing toric CY 4-folds. Namely, usually F-term conditions (defining a master space²) and D-term ones as a whole give a CY 3-fold as the story happens in 4D $\mathcal{N}=1$ superconformal field theories (SCFT). However, because of the appearance of one *dualized photon* $\mathcal{A}$ here, the D-term constraint associated with $\mathcal{A}$ becomes redundant due to *dynamical* Fayet-Iliopoulos (FI) parameters and a 4-fold emerges thereof. Remarkably, we find these properties definitely show up in our model. We demonstrate this mechanism via one explicit example – a 4-fold arising from $\mathbb{C}^3/\mathbb{Z}_3$.

In section 2, a Taub-NUT (ALF) space constructed by means of hyperKähler quotient and its corresponding GLSM are reviewed. Then, in section 3 we write down our model mentioned above whose Higgs branch probes generic 8D hyperKähler manifolds. Section 4 is devoted to the model probing CY 4-folds. Finally, our results are summarized in section 5.

2 Review: 4D Taub-NUT (ALF) space

To begin with, let us see how a multi-centered Taub-NUT (or ALF) space can be constructed by means of hyperKähler quotient [7, 8]. After then, we go to acquaint ourselves

²See [5] for the origin of this terminology. We thanks Forcella for pointing out this reference.

with the GLSM description of it³ according to [2]. The very quotient procedure is exactly carried out in the Higgs branch of the proposed GLSM at IR. Non-zero hyper-multiplets take their values in a hyperKähler manifold thereof.

As said in section 1, from $\mathbb{H}^n \cong \mathbb{R}^{4n}$ one can construct a hyperKähler manifold $\mathbb{H}^n // \mathbb{G}$ with $\mathbb{G}$ being triholomorphic, i.e. it does not alter the hyperKähler structure. We will mainly adopt the following triholomorphic right multiplication

$$\mathcal{Q} \rightarrow \mathcal{Q}e^{i\theta}, \quad \theta \in (0, 4\pi]$$

whose moment map reads

$$\boldsymbol{\mu} = \mathcal{Q}i\bar{\mathcal{Q}} = 2q\tilde{q}k + (|q|^2 - |\tilde{q}|^2)i, \quad (q, \tilde{q}) \in \mathbb{C}^2. \quad (2.1)$$

In addition, a right multiplication $\mathcal{Q}p$ by a unit quaternion p ($p\bar{p} = 1$) $\in SU(2)$ does not vary the hyperKähler structure I^α of $\mathcal{Q}$, while a left multiplication $p\mathcal{Q}$ which keeps the flat metric $ds^2 = d\mathcal{Q}d\bar{\mathcal{Q}}$ invariant but rotates I^α plays the role of $SU(2)_R$ (R -symmetry) under which $(q, \tilde{q}^\dagger)$ transforms as a doublet. Two operations commute. From now on, we usually do not distinguish between $\boldsymbol{\mu}$ being pure imaginary and $\boldsymbol{r} = \left(\Re(2q\tilde{q}), -\Im(2q\tilde{q}), |q|^2 - |\tilde{q}|^2\right) \in \mathbb{R}^3$ with $|\boldsymbol{r}| = |q|^2 + |\tilde{q}|^2$.

Picking up $m+1$ quaternions $\mathcal{Q}_a$ ($a = 1, \dots, m+1$) parameterizing $\mathbb{H}^m \times \mathbb{H} \cong \mathbb{R}^{4m} \times \mathbb{R}^4$, we want to yield a multi-centered Taub-NUT through dividing $\mathbb{H}^{m+1}$ by the following triholomorphic isometry ($\mathcal{Q} = \mathcal{Q}_{m+1}$):

$$\mathcal{Q}_a \rightarrow \mathcal{Q}_a e^{it}, \quad \mathcal{Q} \rightarrow \mathcal{Q} + \lambda t, \quad \lambda \in \mathbb{R}.$$

There are m corresponding triplet moment maps:

$$\boldsymbol{m}_a = \mathcal{Q}_a i \bar{\mathcal{Q}}_a + \frac{\lambda}{2}(\mathcal{Q} - \bar{\mathcal{Q}}) = \boldsymbol{r}_a + \lambda \boldsymbol{y}, \quad \mathcal{Q} = y + \boldsymbol{y}. \quad (2.2)$$

Put into a Taub-NUT form the metric of $\mathbb{H}^m \times \mathbb{H}$ ($r_a = |\boldsymbol{r}_a|$):

$$ds^2 = \sum_a d\mathcal{Q}_a d\bar{\mathcal{Q}}_a = \sum_a \frac{1}{4} \left(\frac{1}{r_a} d\boldsymbol{r}_a^2 + r_a (d\psi_a + \boldsymbol{\omega}_a d\boldsymbol{r}_a)^2 \right) + dy^2 + d\boldsymbol{y}^2, \\ \nabla \times \boldsymbol{\omega}_a = \nabla \frac{1}{r_a}, \quad \psi_a \in (0, 4\pi]. \quad (2.3)$$

³For a multi-centered Taub-NUT space, the corresponding GLSM is considered in [3].

By fixing the level set at $\mathbf{m}_a = \boldsymbol{\zeta}_a$ such that $\mathbf{r} = -\lambda \mathbf{y} = \mathbf{r}_a - \boldsymbol{\zeta}_a$, it is straightforward to show that (2.3) becomes

$$\begin{aligned} ds^2 &= U d\mathbf{r}^2 + \sum_a \frac{|\mathbf{r} + \boldsymbol{\zeta}_a|}{4} (d\psi_a + \boldsymbol{\omega}_a d\mathbf{r}_a)^2 + dy^2, \\ U_a &= \frac{1}{4|\mathbf{r} + \boldsymbol{\zeta}_a|} + \frac{1}{\lambda^2}, \quad U = \sum_a U_a. \end{aligned} \quad (2.4)$$

As a final step, we need to express (2.4) in terms of $\mathbf{r}$ and a *gauge*-invariant χ under moment maps in (2.2), i.e.

$$\chi = \sum_a \psi_a - \frac{y}{\lambda}.$$

Henceforth, instead of ψ 's we use variables like

$$(p_1, p_2, \dots, p_{m-1}), \quad p_\alpha = \sum_{a=1}^{\alpha} \psi_a.$$

The square $\propto (p_\alpha + \dots)^2$ is completed and then dropped due to its *gauge* variance. Finally, we are left with χ and y . Further taking care of the completion of $(y + \dots)^2$, one arrives at

$$ds_{\text{ALF}}^2 = U d\mathbf{r}^2 + \frac{1}{16} U^{-1} \left(d\chi + \left(\sum_{a=1}^m \boldsymbol{\omega}_a \right) d\mathbf{r} \right)^2$$

where $\boldsymbol{\omega}_a$ is evaluated at $\mathbf{r} + \boldsymbol{\zeta}_a$.

2.1 Gauged linear sigma model

Take as the prototype the simplest one-nut TN_4 . Let us review its GLSM following Harvey and Jensen [2]. Introduce $\mathcal{N}=(4,4)$ superfield conventions necessary for later convenience:

$\mathcal{N}=(4,4)$ vector-multiplet (Σ, Φ)

$(\Sigma = \frac{1}{\sqrt{2}} \bar{D}_+ D_- V$ is an $\mathcal{N}=(2,2)$ twisted chiral superfield)

$\mathcal{N}=(4,4)$ hyper-multiplet $(Q, \tilde{Q})$

$\mathcal{N}=(4,4)$ linear hyper-multiplet (Ψ, P)

The GLSM Lagrangian $\mathcal{L} = \mathcal{L}_D + \mathcal{L}_F + \mathcal{L}_{FI}$ for an one-nut TN_4 consists of

$$\begin{aligned}\mathcal{L}_D &= \int d^4\theta \frac{1}{g^2} \Psi^\dagger \Psi + \frac{1}{2g^2} \left(P^\dagger + P + g^2 \sqrt{2} V \right)^2 + \frac{1}{e^2} \left(-\Sigma^\dagger \Sigma + \Phi^\dagger \Phi \right) + \left(Q^\dagger e^{2V} Q + \tilde{Q}^\dagger e^{-2V} \tilde{Q} \right), \\ \mathcal{L}_F &= \int d\theta^+ d\theta^- \mathcal{W} + c.c., \quad \mathcal{W} = \left(\sqrt{2} \tilde{Q} \Phi Q - \Psi \Phi \right), \\ \mathcal{L}_{FI} &= \frac{1}{\sqrt{2}} \int d\theta^+ d\bar{\theta}^- t \Sigma + \int d\theta^+ d\theta^- s \Phi + c.c.. \end{aligned} \tag{2.5}$$

FI terms are included in $\mathcal{L}_{FI}$ with $(s, t) = (\xi^1 - i\xi^2, \xi^3 + i\theta)$ where superscripts are Cartesian labels. To find out the vacuum, one has to expand (2.5) into (bosonic) component fields⁴. Terms from $\mathcal{L}_D$ in order are listed below (Wess-Zumino gauge of V)

$$\begin{aligned} \text{1st :} & \quad -\frac{1}{2g^2} \left((\partial x^1)^2 + (\partial x^2)^2 \right) \\ \text{2nd :} & \quad -\frac{1}{2g^2} \left(2g^4 (\partial \gamma + A)^2 + (\partial x^3)^2 \right) - g^2 |\sigma|^2 \\ \text{3rd :} & \quad \frac{1}{e^2} \left(\frac{1}{2} F_{01}^2 - |\partial \phi|^2 - |\partial \sigma|^2 \right) \\ \text{4th :} & \quad -|\mathcal{D}q|^2 - |\mathcal{D}\tilde{q}|^2 - 2(|q|^2 + |\tilde{q}|^2)|\sigma|^2 \end{aligned} \tag{2.6}$$

Immediately, a D-term potential (by integrating out D in V)

$$V_D = -\frac{e^2}{2} \left(|q|^2 - |\tilde{q}|^2 - (x^3 - \xi^3) \right)^2 \tag{2.7}$$

and F-term potentials (by integrating out $F_Q, F_{\tilde{Q}}, F_\Phi$ and F_Ψ)

$$\begin{cases} V_\Phi = -\frac{e^2}{2} \left| 2q\tilde{q} - (x^1 - ix^2) + s \right|^2 \\ V_Q + V_{\tilde{Q}} = -2|q|^2|\phi|^2 - 2|\tilde{q}|^2|\phi|^2 \\ V_\Psi = -g^2|\phi|^2 \end{cases} \tag{2.8}$$

follow.

Further by taking an infra-red limit ($e^2 \rightarrow \infty$) which decouples the vector-multiplet kinetic term, we find from (2.6) to (2.8) that the *Coulomb* branch is essentially excluded because of $(\phi, \sigma) \rightarrow 0$ ⁵. We finally obtain the supersymmetric vacuum satisfying

$$|q|^2 - |\tilde{q}|^2 = x^3 - \xi^3, \quad 2q\tilde{q} = x^1 - ix^2 - \xi^1 + i\xi^2 \tag{2.9}$$

⁴Bosonic components of Ψ and P are as follows:

$$\Psi = \frac{1}{\sqrt{2}}(x^1 - ix^2) + \dots, \quad P = \frac{1}{\sqrt{2}}(-x^3 + ig^2\gamma) + \dots, \quad \gamma \sim \gamma + 2\pi.$$

⁵There is indeed a term $\mathcal{L}_{top} \sim \theta F_{01}$ though irrelevant here.

which is just the standard moment map considered in (2.1). We observe that the massive gauge field A_μ (mass square g^2) manages to *eat* (gauge away) one unwanted angular variable variant under (2.9). Given that the metric $|dq|^2 + |d\tilde{q}|^2$ of a quaternion is flat, we express it in a Taub-NUT form with $\varphi \sim \varphi + 2\pi$ denoting the triholomorphic $U(1)$ such that

$$\begin{aligned} & |\mathcal{D}q|^2 + |\mathcal{D}\tilde{q}|^2 \Big|_{vacuum} + \frac{1}{2g^2}(\partial\mathbf{x})^2 + \frac{g^2}{2}(\partial\gamma + A)^2 = \\ & \left(\frac{1}{2g^2} + \frac{1}{4|\mathbf{x} - \boldsymbol{\xi}|} \right) (\partial\mathbf{x})^2 + \frac{|\mathbf{x} - \boldsymbol{\xi}|}{4} (2A + 2\partial\varphi + \boldsymbol{\omega}\partial\mathbf{x})^2 + \frac{g^2}{2}(\partial\gamma + A)^2, \\ & \nabla \times \boldsymbol{\omega} = -\nabla \frac{1}{|\mathbf{x} - \boldsymbol{\xi}|}. \end{aligned} \quad (2.10)$$

Defining

$$U = \frac{1}{g^2} + \frac{1}{2|\mathbf{x} - \boldsymbol{\xi}|}, \quad \chi = 2(\gamma - \varphi), \quad \partial Y = -\frac{1}{\sqrt{2}}g(\partial\gamma + A),$$

one obtains

$$\mathcal{L}_{IR} = -\left(\frac{1}{2g^2} + \frac{1}{4|\mathbf{x} - \boldsymbol{\xi}|} \right) (\partial\mathbf{x})^2 - \frac{|\mathbf{x} - \boldsymbol{\xi}|}{4} \left(\partial\chi - \boldsymbol{\omega}\partial\mathbf{x} + 2\sqrt{2}\frac{\partial Y}{g} \right)^2 - (\partial Y)^2.$$

Further by completing $(\partial Y + \dots)^2$,

$$\mathcal{L}_{IR} = -\frac{1}{2}U(\partial\mathbf{x})^2 - \frac{1}{2}U^{-1}\left(\frac{1}{2}\partial\chi - \frac{1}{2}\boldsymbol{\omega}\partial\mathbf{x} \right)^2. \quad (2.11)$$

Note that a remnant $\propto (\partial Y + \dots)^2$ is gauged away by A_μ . To conclude, we have found that at IR ($e^2 \rightarrow \infty$) the Higgs branch vacuum manifold manifests itself as a hyperKähler quotient space. Integrating out A_μ then results in a NL σ M (2.11) with an explicit TN_1 target metric.

3 8D toric hyperKähler manifold

We are mainly interested in 8D toric hyperKähler manifolds $\mathcal{M}_8 = \mathbb{H}^n // \mathbb{G}$ which can be obtained by hyperKähler quotient of quaternions. While $\mathcal{M}_8 = \text{Cone}(\mathcal{B}_7)$ is expressed as a cone, the base seven-manifold $\mathcal{B}_7$ is tri-Sasakian and Einsteinian with constant sectional curvature $R_{\mu\nu} = 6g_{\mu\nu}$. Recently, this kind of geometry has been studied quite intensively

in the context of AdS/CFT because 11D supergravity solutions of the type $AdS_4 \times \mathcal{B}_7$ are dual to various 3D $\mathcal{N}=3$ SCFTs newly found in [9, 10, 11, 12, 13, 14]⁶.

As mentioned before, $\mathbb{G}$ is part of the triholomorphic isometry of $\mathbb{H}^n$ and preserves its hyperKähler structure. With respect to the remaining triholomorphic T^2 , the metric of $\mathcal{M}_8$ can be put into the following form ($\varphi_i \in (0, 4\pi]$):

$$\begin{cases} ds^2 = \frac{1}{2}U_{ij}d\mathbf{x}_i \cdot d\mathbf{x}_j + \frac{1}{2}U^{ij}(d\varphi_i + A_i)(d\varphi_j + A_j) \\ A_i = d\mathbf{x}_j \cdot \boldsymbol{\omega}_{ji} = dx_j^a \omega_{ji}^a, \quad \partial_{x_j^a} \omega_{ki}^b - \partial_{x_k^b} \omega_{ji}^a = \epsilon^{abc} \partial_{x_j^c} U_{ki} \end{cases} \quad (3.1)$$

where $i, j, k = 1, 2$ and $a, b, c = 1, 2, 3$ (Cartesian label)⁷. Conventionally, $\mathcal{M}_8$ is embedded in M-theory and occupies (345678910), i.e. $\mathbf{x}_1 = (345)$ and $\mathbf{x}_2 = (789)$ while circles (φ_1, φ_2) stands for (x^6, x^{10}) . In fact, this metric is so rigid in the sense that it is fully determined once a proper 2 by 2 symmetric matrix U gets specified [6]. Certainly, adding a constant part U_∞ to U is still a solution. Issues about U_∞ will be addressed below.

In generic non-degenerate cases where the holonomy of $\mathcal{M}_8$ is exactly $sp(2)$ (instead of $sp(1) \times sp(1)$), a fraction 3/16 out of full 32 SUSY remains and there exist three real covariantly constant spinors rotated by $SO(3)_R$ R -symmetry of $\mathcal{M}_8$. Since covariantly constant spinors and the triholomorphic isometry commute, the same amount of SUSY survive the duality map which utilizes the above T^2 , say, “(M-theory/two-torus) $\leftrightarrow$ Type IIB string theory”. Consequently, via the duality chain $\mathcal{M}_8$ gets dual to properly-oriented IIB (p, q) 5-branes attached on a circle $\tilde{x}^6$ (T-dualized x^6). How a reduced amount of holonomy from $sp(2)$ to $sp(1) \times sp(1)$ which results in totally 1/4 SUSY occurs? This becomes possible when $\mathcal{M}_8$ just reduces to two orthogonal Taub-NUT spaces occupying (3456) and (78910), respectively (up to some orbifold identification). There may be two kind of situations. One is when $U = U_\infty + \Delta U$ happens to factorize into a diagonal form. The other lies in zooming in on the *near-horizon* region ($\mathbf{x}_{1,2} \sim 0$) such that U_∞ is gotten rid of and ΔU may get properly diagonalized.

As advertised, some comments follow from the relation

$$\frac{U^{-1}}{\sqrt{\det U^{-1}}} = \frac{1}{\Im \tau} \begin{pmatrix} |\tau|^2 & \Re \tau \\ \Re \tau & 1 \end{pmatrix} \quad (3.2)$$

⁶A subfamily of $\mathcal{M}_8$ called Eschenburg space $\mathbb{H}^3//U(1)$ (up to an orbifold identification) is shown to be dual to new $\mathcal{N}=3$ SCFTs in [14].

⁷Note that U^{ij} is the matrix inverse of $U = U_{ij}$.

where we have dropped the subscript ∞ for asymptotic τ and U at infinity. Though (3.2) simply results from translating the asymptotic metric of T^2 in (3.1) into its complex moduli τ , this correlation does imply an $SL(2, \mathbb{Z})$ covariance of both 5-brane charge vectors and the IIB axio-dilaton $\chi + ie^{-\phi} = \tau_\infty$ ⁸. As a matter of fact, U_∞ plays a role of fixing the orientation of 5-branes and subsequently determines the form of ΔU . Therefore as far as preserving six supercharges is concerned, U_∞ and ΔU are not independent. More precisely, the normalization is that when $U_\infty = \mathbf{1}_{2 \times 2}$ ($e^\phi = g_s = 1$) a $(1, k)$ 5-brane⁹ is placed relative to a $(1, 0)$ 5 (NS5) brane occupying (012345) by an angle $\theta = \arctan k$ uniformly on $(3, 7)$ -, $(4, 8)$ - and $(5, 9)$ -plane. In other words, we have measured θ between two kinds of 5-branes according to [6]

$$\cos \theta = \frac{\mathbf{v}^\dagger U_\infty^{-1} \mathbf{v}'}{\sqrt{\mathbf{v}^2 \mathbf{v}'^2}}, \quad \mathbf{v} = (p, q)^t. \quad (3.3)$$

Combining (3.2) and (3.3), we are able to tell why it is $SL(2, \mathbb{Z})$ rather than $SL(2, \mathbb{R})$ that remains as a symmetry of Type IIB string theory by supersymmetry arguments. Because a correct amount of SUSY should be respected in 11D $\mathbb{R}^{1,2} \times \mathcal{M}_8$ due to its hyperKähler nature, dual IIB (p, q) 5-branes are asked to have specific orientations. That (3.3) is $SL(2, \mathbb{Z})$ -invariant leads to $SL(2, \mathbb{Z})$ -covariant $\mathbf{v}$ and $\chi + ie^{-\phi}$ in order to maintain definite 3/16 SUSY necessarily.

3.1 Gauged linear sigma model

Equipped with the above warm-up, we are ready to write down a GLSM which provides triple moment maps from its D- and F-term conditions. It realizes a 8D hyperKähler metric in its Higgs branch thereof at IR upon integrating out auxiliary fields.

To begin with, $\mathcal{L} = \mathcal{L}_D + \mathcal{L}_F$ where

$$\begin{aligned} \mathcal{L}_D &= \int d^4\theta \, \Psi^\dagger \mathcal{G} \Psi + \frac{1}{2g_A^2} (P_A^\dagger + P_A + g_A^2 \sqrt{2} \mathbf{p} \cdot \mathbf{V})^2 + \frac{1}{2g_B^2} (P_B^\dagger + P_B + g_B^2 \sqrt{2} \mathbf{q} \cdot \mathbf{V})^2 \\ &+ \sum_{i=1}^M \left(\frac{1}{e_i^2} (-\Sigma_i^\dagger \Sigma_i + \Phi_i^\dagger \Phi_i) + H_i^\dagger e^{2V_i} H_i + \tilde{H}_i^\dagger e^{-2V_i} \tilde{H}_i \right), \\ \mathcal{L}_F &= \int d\theta^+ d\theta^- \, \mathcal{W} + c.c., \quad \mathcal{W} = \left(\mathbf{H} \tilde{\mathbf{H}} \cdot \boldsymbol{\Phi} \right) - \left(\Psi_A \mathbf{p} + \Psi_B \mathbf{q} \right) \cdot \boldsymbol{\Phi}. \end{aligned} \quad (3.4)$$

⁸As is well-known, $\chi + ie^{-\phi}$ gets identified with the M-theory torus's complex moduli τ_∞ via the aforementioned duality chain.

⁹It occupies $012(3, 7)_\chi(4, 8)_\psi(5, 9)_\theta$.

This is an extension of the original work of Okuyama [3]. Our convention is as follows:

$$\begin{aligned}\Psi^\dagger &= (\Psi_A, \Psi_B), \quad \mathcal{G} = \begin{pmatrix} \frac{1}{g_A^2} & 0 \\ 0 & \frac{1}{g_B^2} \end{pmatrix}, \\ \Psi_{A,B} &= \frac{1}{\sqrt{2}}(x_{A,B}^1 - ix_{A,B}^2) + \cdots, \quad P_{A,B} = \frac{1}{\sqrt{2}}(-x_{A,B}^3 + ig^2\gamma_{A,B}) + \cdots, \quad \gamma_{A,B} \sim \gamma_{A,B} + 2\pi.\end{aligned}\tag{3.5}$$

Bold symbols (e.g. $\mathbf{p}, \mathbf{q}, \Phi, \mathbf{V}, \mathbf{H}\tilde{\mathbf{H}}$, etc.) denote M -vectors, while $\cdot$ represents the usual vector inner product. Also, $\mathcal{G}$ will be identified with the U_∞ part in (3.1). We have turned off FI parameters which spoil the cone structure of $\mathcal{M}_8$ when one zooms in on the vicinity around the origin. In addition to couplings constants, $(\mathbf{p}, \mathbf{q})$ are the only parameters in (3.4). It becomes transparent later that $(\mathbf{p}, \mathbf{q})$ is responsible for M distinct charge vectors of IIB (p, q) 5-branes.

We should expand $\mathcal{L}$ into components fields, integrate out auxiliary D and F fields. Further imposing the infra-red limit $e_i^2 \rightarrow \infty$ freezes kinetic terms of vector-multiplets. Henceforth, we are left with (bosonic part only)

$$\begin{aligned}& -\frac{1}{2g_A^2}(\partial\mathbf{x}_A)^2 - \frac{1}{2g_B^2}(\partial\mathbf{x}_B)^2 - g_A^2(\partial\gamma_A + \mathbf{p} \cdot \mathbf{A})^2 - g_B^2(\partial\gamma_B + \mathbf{q} \cdot \mathbf{A})^2 \\ & + \sum_i \left(-g_A^2|p_i\sigma_i|^2 - g_B^2|q_i\sigma_i|^2 - |\mathcal{D}h_i|^2 - |\mathcal{D}\tilde{h}_i|^2 - 2(|h_i|^2 + |\tilde{h}_i|^2)|\sigma_i|^2 \right)\end{aligned}\tag{3.6}$$

and potentials

$$\begin{cases} V_D = \sum_i -\frac{e_i^2}{2} \left(|h_i|^2 - |\tilde{h}_i|^2 - (p_i x_A^3 + q_i x_B^3) \right)^2 \\ V_F = \sum_i -\frac{e_i^2}{2} \left| 2h_i\tilde{h}_i - p_i(x_A^1 - ix_A^2) - q_i(x_B^1 - ix_B^2) \right|^2 - |\phi_i|^2(2|h_i|^2 + 2|\tilde{h}_i|^2 + g_A^2 p_i^2 + g_B^2 q_i^2) \end{cases}$$

A SUSY vacuum accompanied by $(\phi_i, \sigma_i) \rightarrow 0$ implies M sets of triple moment maps:

$$\begin{cases} |h_i|^2 - |\tilde{h}_i|^2 - (p_i x_A^3 + q_i x_B^3) = 0 \\ 2h_i\tilde{h}_i - p_i(x_A^1 - ix_A^2) - q_i(x_B^1 - ix_B^2) = 0 \end{cases}\tag{3.7}$$

from $V_D = V_F = 0$. It is now clear that the model itself performs a hyperKähler quotient on $M + 2$ quaternions by imposing (3.7) which kill M pairs of $(h_i, \tilde{h}_i)$.

As before, let us rewrite the flat metric $|dh|^2 + |d\tilde{h}|^2$ of a quaternion into a Taub-NUT

form with ψ denoting the triholomorphic $U(1)$, i.e.

$$\begin{aligned} & \sum_{i=1}^M |\mathcal{D}h_i|^2 + |\mathcal{D}\tilde{h}_i|^2 \Big|_{vacuum} \\ &= \Delta U_{ab} \partial \mathbf{x}_a \cdot \partial \mathbf{x}_b + \sum_{i=1}^M \frac{|\mathbf{X}_i|}{4} (2A_i + 2\partial\psi_i + \boldsymbol{\omega}_i \cdot \partial \mathbf{X}_i)^2, \quad a, b = A, B \end{aligned}$$

where

$$\Delta U = \sum_{i=1}^M \frac{1}{4|\mathbf{X}_i|} \begin{pmatrix} p_i^2 & p_i q_i \\ p_i q_i & q_i^2 \end{pmatrix}, \quad \mathbf{X}_i = p_i \mathbf{x}_A + q_i \mathbf{x}_B, \quad \nabla \times \boldsymbol{\omega}_i = -\nabla \frac{1}{|\mathbf{X}_i|}.$$

According to [6], ΔU corresponds to M dual IIB (p_i, q_i) 5-branes localized at $\mathbf{X}_i = 0$ w.r.t. $U_\infty = \mathcal{G}$ which can be adjusted to $(\text{const.})\mathbf{1}_{2 \times 2}$. Since only $\chi_A = \gamma_A - \sum p_i \psi_i$ and $\chi_B = \gamma_B - \sum q_i \psi_i$ are invariant under the action of moment maps in (3.7), our strategy is to let M massive gauge fields gauge away remaining *gauge*-variant angular variables. To evaluate the term

$$\sum_{i=1}^M \frac{|\mathbf{X}_i|}{4} (2A_i + 2\partial\psi_i + \boldsymbol{\omega}_i \cdot \partial \mathbf{X}_i)^2 - \frac{g_A^2}{2} (\partial\gamma_A + \mathbf{p} \cdot \mathbf{A})^2 - \frac{g_B^2}{2} (\partial\gamma_B + \mathbf{q} \cdot \mathbf{A})^2, \quad (3.8)$$

we adopt variables

$$\tau_k = \sum_{i=1}^k p_i \mathcal{B}_i, \quad \tilde{\tau}_k = \sum_{i=1}^k q_i \mathcal{B}_i, \quad \mathcal{B}_i = A_i + \partial\psi_i, \quad k = 1, \dots, M$$

such that every $2(A_i + \partial\psi_i)$ splits into a sum

$$\frac{\tau_i - \tau_{i-1}}{p_i} + \frac{\tilde{\tau}_i - \tilde{\tau}_{i-1}}{q_i}.$$

In dealing with (3.8), we complete in order $(\zeta_k \tau_k + \eta_k \tilde{\tau}_k + \dots)^2$ ($\zeta_k, \eta_k = \text{const.}$) from $k = 1$ and gauge them away. Finally by substituting the outcome into (3.6) the desired metric form as (3.1) can be reached, i.e.

$$\begin{aligned} \mathcal{L}_{IR} &= -U_{ab} \partial \mathbf{x}_a \cdot \partial \mathbf{x}_b - \frac{1}{4} U^{ab} (\partial\chi_a + \mathcal{E}_a) (\partial\chi_b + \mathcal{E}_b), \quad \chi_a \sim \chi_a + 2\pi, \\ \mathcal{E}_a &= -\frac{1}{2} \sum_{i=1}^M v_{i,a} \boldsymbol{\omega}_i \cdot \partial \mathbf{X}_i, \quad v_i = (p_i, q_i). \end{aligned}$$

This is exactly a NL σ M which pulls back the metric of $\mathcal{M}_8$.

4 $\mathcal{N}=(2,2)$ GLSM for Calabi-Yau fourfold

Let us propose an $\mathcal{N}=(2,2)$ GLSM whose IR moduli space becomes a CY 4-fold. Our superfield conventions are inherited from section 3. The Lagrangian under consideration is $\mathcal{L} = \mathcal{L}_D + \mathcal{L}_F$ where¹⁰

$$\begin{aligned}\mathcal{L}_D &= \int d^4\theta \frac{1}{2g^2} (P^\dagger + P + \mathbf{k} \cdot g^2 \sqrt{2} \mathbf{V})^2 - \sum_{a=1}^r \frac{1}{e_a^2} \Sigma_a^\dagger \Sigma_a \\ &+ \sum_i \mathcal{H}_i^\dagger e^{2V_{h(i)}} \mathcal{H}_i e^{-2V_{t(i)}} + \sum_{a=1}^r \Phi_a^\dagger e^{2V_a} \Phi_a, \\ \mathcal{L}_F &= \int d\theta^+ d\theta^- \mathcal{W} + c.c., \quad \mathbf{k} = (k_1, \dots, k_r) \in \mathbb{Z}^r.\end{aligned}\tag{4.1}$$

Note that $\mathcal{W}$ denotes the superpotential and the gauge invariance in the first term of $\mathcal{L}_D$ is maintained by

$$\mathbf{k} \cdot \mathbf{V} \rightarrow \mathbf{k} \cdot \mathbf{V} + (\Lambda^\dagger + \Lambda), \quad P \rightarrow P - \sqrt{2}g^2\Lambda, \quad P^\dagger \rightarrow P^\dagger - \sqrt{2}g^2\Lambda^\dagger.$$

Bi-fundamental chiral fields $\mathcal{H}_i$ charged under r $U(1)$'s according to a given quiver diagram with r nodes. The subscript i runs over arrows representing bi-fundamental fields in the quiver diagram, while $h(i)$ or $t(i)$ specifies a node on which the head or tail of i ends. In addition, $\mathcal{W}$ subject to holomorphy is again related to the given quiver¹¹. Note that only P of the previous $\mathcal{N}=(4,4)$ multiplet (Ψ, P) enters $\mathcal{L}$ with the lowest component $x + i\gamma$ ($\gamma \sim \gamma + 2\pi$). It is apparent that both $\mathcal{L}$ and the resultant IR moduli space are characterized only by $\mathbf{k}$ and the responsible quiver diagram.

First of all, let us expand superfields in $\mathcal{L}$ into components and integrate out auxiliary D and F fields. Taking $(e_a^2, g^2) \rightarrow \infty$ freezes kinetic terms of fields (A_a, σ_a, x) . What we are left with are (bosonic part only)

1. Kinetic terms of (h_i, ϕ_a) (bosonic component of $(\mathcal{H}_i, \Phi_a)$)
2. A remnant:

$$-g^2(\partial\gamma + \mathbf{k} \cdot \mathbf{A})^2\tag{4.2}$$

¹⁰We noticed incidentally during revising this version that there is a similar D-term in four dimensions considered in the context of “chaotic D-term inflation” by Kawano [15].

¹¹Based on the *brane tiling* technology one can read off $\mathcal{W}$ from a given quiver. See [16] for an excellent review.

3. A potential:

$$V_{pot} = \sum_i |h_i|^2 |Q_i^{h(i)} \sigma_{h(i)} - Q_i^{t(i)} \sigma_{t(i)}|^2 + \sum_a 2|\sigma_a|^2 |\phi_a|^2 \\ + \sum_a g^2 |k_a \sigma_a|^2 + \sum_i \left| \frac{\partial \mathcal{W}}{\partial h_i} \right|^2 + \sum_a \left| \frac{\partial \mathcal{W}}{\partial \phi_a} \right|^2 + V_D,$$

where

$$V_D = \sum_a \frac{e_a^2}{2} \left(\sum_i (\delta_{h(i)}^a Q_i^a - \delta_{t(i)}^a Q_i^a) |h_i|^2 + |\phi_a|^2 - k_a x \right)^2 = \sum_a \frac{e_a^2}{2} D_a^2 \quad (4.3)$$

The vacuum manifold $\mathcal{M}$ is determined by

$$\sigma_a = \phi_a = 0 \quad D_a = 0, \quad d\mathcal{W} = 0. \quad (4.4)$$

Several comments follow. A distinguishing feature departing from those dealt with in previous sections is that by means of γ 's e.o.m. one gauge field $\mathcal{A} = \mathbf{k} \cdot \mathbf{A}$ in (4.2) can be dualized into a scalar $-\gamma$. In the context of 3D Yang-Mills-Chern-Simons theory, $\mathcal{A}$ is referred to as a *dual photon*. Together with $V_D = 0$ we can understand that the model itself executes rather a *symplectic* quotient over complex variables h_i than a hyperKähler one.

Let us analyze the moduli space in detail from both field-theoretical and geometrical grounds.

1. The constraint $\sum_a D_a = 0$ in (4.3) characterizes the decoupled diagonal $U(1)$ and implies as well that $\sum_a k_a = 0$ or the vector $\mathbf{k}$ is orthogonal to $(1, \dots, 1)$. To this end, one has only $(r-2)$ linearly independent D-term conditions [17]:

$$\sum_a \ell_a D_a = 0, \quad \ell \in \text{Ker}(\mathbf{k}), \quad \ell \neq (1, \dots, 1).$$

This fact is consistent with that we have a dualized $\mathcal{A}$ and leads naturally to $\mathcal{M} = \{d\mathcal{W} = 0\} // C^{\star r-2}$ which is a CY 4-fold by definition emerging from a 3-fold. Namely, $\mathcal{M} = \{d\mathcal{W} = 0\} // C^{\star r-1}$ is a CY 3-fold because its derivation is just the same with that of 4D $\mathcal{N}=1$ SCFTs on a bunch of D3-branes probing CY₃ cones.

2. Geometrically, an ungauged $U(1)$ $\mathcal{A}$ suggests the existence of an S^1 in the vacuum

moduli space correspondingly. According to $V_D = 0$, we can thus describe $\mathcal{M}$ as a 3-fold fibered on a real line x with a circle bundle fibered over them with non-trivial first Chern class [18, 19]. We find that the field x in (4.3) (or $V_D = 0$) serves as *dynamical* FI parameters and plays the very role of giving varying Kähler classes for the fibered 3-fold.

3. Let us elaborate on arguments about the *surviving* S^1 bundle associated with $\mathcal{A}$. For conceptual convenience, we can first think of it as γ charged logarithmically under r $U(1)$'s:

$$\gamma \rightarrow \gamma + \sum_{a=1}^r k_a \lambda_a, \quad \gamma \sim \gamma + 2\pi. \quad (4.5)$$

Consequently, (4.5) defines a one-parameter gauge transformation $\boldsymbol{\lambda} = t(k_1, \dots, k_r)$. We find that γ is especially helpful when one wants to determine the correct periodicity of S^1 . In fact, there may remain some discrete amount of gauge symmetry Γ of $\mathcal{A}$ such that Γ shrinks the circumference of S^1 to $2\pi/\gcd(\mathbf{k})$. This is apparent from the expression of $\boldsymbol{\lambda}$. Let us put things in an inverse way. From (4.2) $\mathcal{A}$ is spontaneously broken once γ acquires a vev γ_0 . But if there exists $f = \gcd(\mathbf{k})$ with $\gamma_0 = f\gamma'_0 + 2m\pi$ ($m \in \mathbb{Z}$), the gauge fixing becomes incomplete due to a factor $2m\pi/f$. This soon means that the gauge symmetry of $\mathcal{A}$ is only spontaneously broken down to a discrete extent. To conclude, we should at the end divide $\mathcal{M}$ further by Γ . All these are directly reminiscent of the novel mechanism in newly-developed 3D $\mathcal{N}=2$ Chern-Simons-matter theories [20]. Therefore, it turns out that, up to a discrete quotient Γ , $\mathcal{M} = \{d\mathcal{W} = 0\}/C^{\star r-2}$ where additional real two dimensions result effectively from degrees of freedom of $x + i\gamma \subset P^{12}$.

4.1 Toric geometry

Let us review quickly stuffs about toric varieties which are highly helpful when one lifts a toric CY_3 to a toric CY_4 . In general, a toric variety V is expressed as $\mathbb{C}^n \setminus F // \mathbb{C}^{\star \mathcal{G}}$ and can be summarized pictorially by a toric diagram¹³ consisting of n d -vectors $\nu_i \in \mathbb{Z}^d$ ($d = n - \mathcal{G}$) subject to

$$\sum_{i=1}^n Q_i^A \nu_i = 0, \quad A = 1, \dots, \mathcal{G}, \quad Q_i^A : \text{charge matrix}. \quad (4.6)$$

¹²Typically, x and γ are called Fayet-Iliopoulos and Stueckelburg fields, respectively.

¹³The subset F is determined by partial resolutions.

By imposing the Calabi-Yau condition: $\sum_i Q_i^A = 0 \quad \forall A$, these ν 's can be written as $\nu_i = (1, \dots)^t$ and are called toric data of a toric CY d -fold. For $d=3$, toric data are plotted on a plane, while $d=4$ toric data are encoded in 3D lattice points which define a convex polytope – *crystal*. This is why we adopt the name *crystal* in our title page because CY 4-folds are under consideration.

Let us talk about the geometric meaning of toric data. Every ν_i assigns a shrinking 1-cycle out of T^d at the i -th facet of the boundary $\partial\mathcal{C}_+$ where $\mathcal{C}_+$ denotes a cone in $\mathbb{Z}^d$ over which T^d is fibered over. In $d=3$ cases, lines where two facets meet will altogether constitute a *web* diagram which in turn represents a tree formed by multiple semi-infinite NS5-branes. Although we will not go further to details, interested readers are encouraged to consult [16].

4.2 Calabi-Yau crystal

Though lifting 3-folds to 4-folds has been studied quite heavily to date¹⁴, let us just pick up one canonical example $\mathbb{C}^3/\mathbb{Z}_3$ discussed also in [17, 24, 19]. Here, $\mathbb{Z}_3$ is embedded in the Hopf $U(1)$ bundle of $\mathbb{C}^3$ whose action is

$$(z_1, z_2, z_3) \rightarrow (\epsilon z_1, \epsilon z_2, \epsilon z_3), \quad \epsilon = e^{2\pi i/3}. \quad (4.7)$$

4.2.1 $\mathbb{C}^3/\mathbb{Z}_3$

A stack of D3-branes probing $\mathbb{C}^3/\mathbb{Z}_3$ [25, 26] has nine bi-fundamental chiral superfields on its worldvolume gauge theory with a superpotential (though we will focus on its abelian version)

$$\mathcal{W} \propto \epsilon^{IJK} \text{Tr}(X_I Y_J Z_K), \quad I, J, K = 1, 2, 3. \quad (4.8)$$

In Figure 1, three nodes denote three gauge groups and hence (3–1) D-term conditions. In addition, $d\mathcal{W} = 0$ renders four more constraints

$$X_1 Y_2 = X_2 Y_1, \quad X_2 Y_3 = X_3 Y_2, \quad (X, Y) \leftrightarrow (Y, Z). \quad (4.9)$$

¹⁴See also [21, 22, 23] and references therein.




Figure 1: A quiver diagram for $\mathbb{C}^3/\mathbb{Z}_3$. Three nodes are labeled by integers (k_1, k_2, k_3) . Three sets of chiral superfields are denoted as (X_I, Y_I, Z_I) ($I = 1, 2, 3$).

Instead, we can *solve* (4.9) via $X_I = u_1 v_I$, $Y_I = u_2 v_I$ and $Z_I = u_3 v_I$ ($I = 1, 2, 3$) accompanied by a D-term like condition:

$$|u_1|^2 + |u_2|^2 + |u_3|^2 - |v_1|^2 - |v_2|^2 - |v_3|^2 = 0. \quad (4.10)$$

$\mathcal{M}_F = \{d\mathcal{W} = 0\}$ can then be thought of as $\mathbb{C}^6//\mathbb{C}^\star_{(1,1,1,-1,-1,-1)}$. Alternatively, the following exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{Q^t} \mathbb{Z}^6 \xrightarrow{T} \mathbb{N} \rightarrow 0 \quad (4.11)$$

reveals the same thing. It merely says mathematically $\mathcal{M}_F$ can be regarded as $\mathbb{C}^6//\mathbb{C}^\star$ where the action of $\mathbb{C}^\star$ is spelt out by Q . More precisely, one can express $\mathcal{M}_F$ as a cone $\mathbb{M}_+$ in $\mathbb{M} = \mathbb{Z}^6$. Its dual cone (toric data) $\mathbb{N}_+$ in $\mathbb{N} = \text{Hom}(\mathbb{M}, \mathbb{Z})$ is generated by six lattice points, i.e. $T: \mathbb{Z}^6 \rightarrow \mathbb{N}$. By definition $\text{Ker}(T)$ is the image of Q^t , so the cokernel of Q^t is isomorphic to T (Q : 1 by 6 matrix). To get a 3-fold, we need to incorporate D-term constraints as well. In summary, we can obey three steps below [26]:

1. Introduce U (5 by 6 matrix) with $UT^t = 1$ and V (2 by 5 matrix). V encodes two D-term constraints on five independent variables as indicated in (4.9)

2. Concatenate VU (2 by 6 matrix) and Q to yield Q' (3 by 6 matrix)
3. Computing the cokernel of Q'^t gives us desired toric data in terms of a map $\mathcal{T}: \mathbb{Z}^6 \rightarrow \mathbb{Z}^3$

To have a 4-fold we need to use a new V involving only one linearly-combined D-term constraint (irrelevant to $\mathcal{A}$) out of three nodes and repeat the above procedure. Instead of this approach, a more convenient one figured out in [17] is to directly perform one more quotient on previous (u_I, v_I) by another $\mathbb{C}^\star$: $(k_1 + k_2, -k_1, -k_2, 0, 0, 0)$. Translating this into the action on (X_I, Y_I) , one easily realizes that it is the combination $-k_2 Q_1 + k_1 Q_2$ orthogonal to $\mathbf{k} \cdot \mathbf{Q}$ with $\mathbf{k} = (k_1, k_2, k_3 = -k_1 - k_2)$. See Table 1 for informations about Q_i responsible for the i -th node. Arbitrary $\mathbf{k}$ encountered in (4.1) meet the requirement

Table 1: Nine bi-fundamental chiral matters charged under three gauge groups in the case of $\mathbb{C}^3/\mathbb{Z}_3$

	X_1	X_2	X_3	Y_1	Y_2	Y_3	Z_1	Z_2	Z_3
Q_1	-1	-1	-1	0	0	0	1	1	1
Q_2	1	1	1	-1	-1	-1	0	0	0
Q_3	0	0	0	1	1	1	-1	-1	-1

$\sum_a k_a = 0$. Also, we may demand $\gcd(k_1, k_2) = 1$ technically in order to avoid a further discrete quotient Γ explained around (4.5).

All in all, via $(k_1, k_2) = (2p - k, k - p)$ we end up with an eight cone over a seven Sasaki-Einstein base denoted as $Y^{p,k}(\mathbb{CP}^2)$. Its toric data are [17]:

$$\begin{aligned}
w_1 &= (0, 0, 0), & w_2 &= (0, 0, p), & w_3 &= (1, 0, 0), \\
w_4 &= (0, 1, 0), & w_5 &= (-1, -1, k), & w_0 &= (0, 0, k - p).
\end{aligned} \tag{4.12}$$

These six vectors do form a 3D crystal as was shown in [27] with an additional vector w_0 standing for a resolution, provided $p \leq k \leq 2p$. See [17] for more details.

5 Summary

In this short note we have studied two kinds of GLSMs whose vacuum moduli spaces are real eight-dimensional hyperKähler and Calabi-Yau manifolds, respectively.

The Higgs branch of the former $\mathcal{N}=(4,4)$ GLSM manifests itself as a generic 8D hyperKähler quotient space at the infra-red regime. On the other hand, the latter $\mathcal{N}=(2,2)$ GLSM has a more flexible superpotential $\mathcal{W}$ as well as chiral matter contents spelt out by a quiver diagram. Usually, D- and F-term constraints give rise to a toric CY 3-fold just as in familiar 4D $\mathcal{N}=1$ SCFT setups. Nevertheless, owing to a novel mechanism introduced in section 4, 3-folds are promoted to 4-folds exactly the same way as what happens in recently-developed 3D $\mathcal{N}=2$ Chern-Simons-matter theories which have toric CY 4-folds as their moduli spaces. In words, in addition to the diagonal gauge field $\sum_a A_a$, an ungauged dual photon $\mathcal{A} = \sum_a k_a A_a$ as well does not enter the quotient process due to *dynamical* Fayet-Iliopoulos parameters. Therefore, the moduli space dimension is enhanced from six to eight. The extra two dimensions effectively result from degrees of freedom of a chiral superfield P . Parameters k_a in our 2D Lagrangian satisfying $\sum_a k_a = 0$ are arbitrary integers. Their role is analogous to 3D quiver Chern-Simons levels. We discussed the whole story via a very well-developed 3-fold $\mathbb{C}^3/\mathbb{Z}_3$.

Acknowledgements

We would like to thank Amihay Hanany, Kazuo Hosomichi, Hiroaki Kanno, Muneto Nitta, Hiroshi Ohki, Yutaka Sakamura and especially Kazumi Okuyama for enlightening discussions at different intermediate stages. TST is grateful to organizers of the wonderful workshop on “Branes, Strings and Black Holes” at Yukawa Institute of Theoretical Physics. TST is supported in part by the postdoctoral program at RIKEN.

References

- [1] E. Witten, Nucl. Phys. B **403**, 159 (1993) [arXiv:hep-th/9301042].
- [2] J. A. Harvey and S. Jensen, JHEP **0510** (2005) 028 [arXiv:hep-th/0507204].
- [3] K. Okuyama, JHEP **0508** (2005) 089 [arXiv:hep-th/0508097].
- [4] K. Okuyama, Phys. Lett. B **668** (2008) 153 [arXiv:0807.0047 [hep-th]].
- [5] D. Forcella, A. Hanany, Y. H. He and A. Zaffaroni, JHEP **0808** (2008) 012 [arXiv:0801.1585 [hep-th]].

- [6] J. P. Gauntlett, G. W. Gibbons, G. Papadopoulos and P. K. Townsend, Nucl. Phys. B **500** (1997) 133 [arXiv:hep-th/9702202].
- [7] G. W. Gibbons and P. Rychenkova, Commun. Math. Phys. **186** (1997) 585 [arXiv:hep-th/9608085].
- [8] E. Witten, JHEP **0906** (2009) 067 [arXiv:0902.0948 [hep-th]].
- [9] Y. Imamura and K. Kimura, Prog. Theor. Phys. **120** (2008) 509 [arXiv:0806.3727 [hep-th]].
- [10] D. L. Jafferis and A. Tomasiello, JHEP **0810** (2008) 101 [arXiv:0808.0864 [hep-th]].
- [11] S. Hohenegger and I. Kirsch, JHEP **0904** (2009) 129 [arXiv:0903.1730 [hep-th]].
- [12] D. Gaiotto and D. L. Jafferis, “Notes on adding D6 branes wrapping $\mathbb{RP}^3$ in $AdS_4 \times \mathbb{CP}^3$,” arXiv:0903.2175 [hep-th].
- [13] Y. Hikida, W. Li and T. Takayanagi, JHEP **0907** (2009) 065 [arXiv:0903.2194 [hep-th]].
- [14] M. Fujita and T. S. Tai, JHEP **0909** (2009) 062 [arXiv:0906.0253 [hep-th]].
- [15] T. Kawano, Prog. Theor. Phys. **120** (2008) 793 [arXiv:0712.2351 [hep-th]].
- [16] M. Yamazaki, Fortsch. Phys. **56** (2008) 555 [arXiv:0803.4474 [hep-th]].
- [17] D. Martelli and J. Sparks, Phys. Rev. D **78** (2008) 126005 [arXiv:0808.0912 [hep-th]].
- [18] M. Aganagic and C. Vafa, arXiv:hep-th/0110171.
- [19] M. Aganagic, arXiv:0905.3415 [hep-th].
- [20] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, JHEP **0810** (2008) 091 [arXiv:0806.1218 [hep-th]].
- [21] K. Ueda and M. Yamazaki, JHEP **0812** (2008) 045 [arXiv:0808.3768 [hep-th]].
- [22] Y. Imamura and K. Kimura, JHEP **0810** (2008) 114 [arXiv:0808.4155 [hep-th]].
- [23] M. Taki, arXiv:0910.0370 [hep-th].
- [24] A. Hanany and A. Zaffaroni, JHEP **0810** (2008) 111 [arXiv:0808.1244 [hep-th]].
- [25] M. R. Douglas and G. W. Moore, arXiv:hep-th/9603167.
- [26] M. R. Douglas, B. R. Greene and D. R. Morrison, Nucl. Phys. B **506** (1997) 84 [arXiv:hep-th/9704151].

[27] D. Martelli and J. Sparks, JHEP **0811** (2008) 016 [arXiv:0808.0904 [hep-th]].